

APPLICATION UNDER UNITED STATES PATENT LAWS

Atty, Dkt. No. PW 282799
(M#)

Invention: PEPTIDES CAPABLE OF FUNCTIONING AS MIMOTOPES FOR HORMONAL ANALYTES

Inventor (s): BADLEY, Robert A
BERRY, Mark J
WILLIAMS, Samantha C

Pillsbury Winthrop LLP
Intellectual Property Group
1600 Tysons Boulevard

McLean, VA 22102
Attorneys
Telephone: (703) 905-2000

This is a:

- ☐ Provisional Application
- ☒ Regular Utility Application
- ☐ Continuing Application
 - ☒ The contents of the parent are incorporated by reference
- ☐ PCT National Phase Application
- ☐ Design Application
- ☐ Reissue Application
- ☐ Plant Application
- ☐ Substitute Specification
 - Sub. Spec Filed _____
 - in App. No. _____ / _____
- ☐ Marked up Specification re
 - Sub. Spec. filed _____
 - In App. No. _____ / _____

SPECIFICATION

Title: Peptides Capable of Functioning as Mimotopes for Hormonal Analytes

Field of the Invention

This invention relates to the discovery that certain peptide molecules have similar reactive properties as certain steroidal compounds, notwithstanding the significant structural dissimilarities between such compounds, and are thus capable of functioning as mimotopes of the steroidal compounds in, for example, displacement immunoassays designed for the detection of steroids.

Background of the Invention

As a matter of general definition, an epitope is that region of a particular antigen which contains the critical binding region of the antigen necessary for triggering an immunity-related antibody binding response. Epitopes are also often referred to in the alternative as antigenic determinants.

Understanding the structures of epitopes as well as their specific binding reactions to particular antibodies is of significant interest to many, as such an understanding could lay the foundation for advancements in the pharmaceutical, diagnostic and health industries. To facilitate this understanding, in recent years academic institutions and industry have constructed what are termed epitope libraries.

Epitope libraries are large collections of variable amino acid sequences that are displayed, for example, on the surfaces of bacteriophage. Each sequence corresponds to a particular epitope of a particular antigen. Often, the epitope libraries will consist of many millions of these short amino acid sequences, sometimes even as many as one hundred million sequences or more. Representative epitope libraries are described in detail in Luzzago A., et al. *Mimicking of discontinuous epitopes by phage-displayed peptides, I. Epitope mapping of human H ferritin using a phage library of constrained peptides*. Gene 128, 51-57 (1993).

Once an epitope library is constructed, antibodies or other binding proteins can be utilised to select specifically for a particular epitope. The epitope can then be sequenced, either directly or by first identifying the corresponding DNA sequence and then by transcribing and translating that DNA sequence into the corresponding amino acid sequence. By such techniques, the binding regions of antigenic compounds and molecules can be determined; and it can be readily envisioned that once the binding regions of particular antigens are known, powerful biotechnology applications -- such as the design of vaccines using particular epitopes -- can be achieved.

Despite the apparent power of epitope library screening techniques, they have heretofore been used primarily only to identify and sequence the specific binding regions of particular antigens. This in and of itself serves to limit the type and scope of biotechnology applications that can be based on this technology.

In Scott, *Discovering Peptide Ligands Using Epitope Libraries*, Trends in Biochemical Science 17, pp. 241-245 (July 1992), an extension of conventional epitope library techniques is disclosed. Specifically, Scott asserts that epitope libraries can be used to identify and map peptide mimotopes for known antigens. A mimotope is a molecular sequence which "mimics" the epitopic region of a particular antigen, but which does not contain the specific amino acid sequence which comprises the epitope. Thus, a mimotope is structurally distinct from an epitope, though functionally it is very similar as it is capable of binding in a similar fashion to the binding cleft of the antibody directed to the antigen containing the particular epitope.

Though mimotopes technically can be any molecules or sequence of molecules which mimic an epitope, they are most often small, low molecular weight peptides which comprise short sequences of amino acids. Because they are most typically small peptides, they have been thought to be constrained in what they can mimic. Specifically, it has been a generally held belief that peptide mimotopes could be identified for numerous protein based antigens (i.e. those antigens with peptide epitopes), but that because of the complex structure of antibody binding clefts, and the correspondingly complex nature of the

antibody binding response, all of which it was believed would require a close similarity in structure between the mimotope and epitope in order for mimotope-antibody binding to occur, the identification of peptide mimotopes for non-protein based antigens would be difficult to achieve. In general this belief has been proved correct, although there are a few isolated exceptions.

For example, in *Random peptide libraries: A source of specific binding molecules*: Devlin JJ: Science 249, 404-406 (1990), peptide mimotopes have been identified for biotin, an essential vitamin necessary for certain enzymatic carboxylation reactions in living cells. Though biotin is not peptidal in structure, it is nevertheless similar in size and structure to several amino acids (for example, histidine). Thus, it was not unexpected that a peptide mimotope for such a molecule could be identified.

Similarly, in *Peptide ligands for a sugar binding protein isolated from a random peptide library*: Oldenburg, KR et al., Proc. Nat. Acad. Sci. USA, 89, 5393-5397 (1992), peptide mimotopes for the mannopyranoside ligand of concanavalin A have been identified. Again though, such mimotopes are of similar size and have a similar structural configuration to the epitopes which they mimic and thus their identification was less than surprising. Mimotopes of certain forms of DNA have also been identified, as described in Sibille et al., *Mimotopes of polyreactive anti-DNA antibodies identified using phage displayed peptide libraries*: Eur J Immunol., 27, 1221-1228 (1997)

Notwithstanding these few limited discoveries, the identification of peptide mimotopes from epitope libraries (or other means) for complex non-protein molecules such as steroidal compounds has not occurred. Furthermore, due to the clear structural differences between such peptide mimotopes and complex non-protein molecules such as steroids, the mere existence of these mimotopes has heretofore been questioned.

WO 96/16322 discloses an affinity-based process for recovering specific binding agents with high affinity for a particular target ligand, which involves the use of first and second analogues of the target ligand. Page 4 of the document mentions that one of the analogues

used in the process may be an epitope mimic, "i.e. a small molecule, generally of synthetic origin, such as a short peptide, which behaves in a manner comparable to the binding site (epitope) of the target ligand".

Although the document includes examples of the process in which the target ligand is a steroid (specifically, estrone-3-glucuronide, "E3G") none of these examples involve the use of a peptide mimotope of a steroid, and there is no disclosure of a specific example of, nor any experimental evidence relating to, a peptide mimotope of a steroid. Indeed at page 4, immediately following the passage quoted above, WO 96/16322 states "*Epecially when the target ligand is E3G, said first analogue can be estrone. Preferably said second analogue is estriol glucuronide. Alternatively, estradiol-3-glucuronide can be used as the second analogue; in this case, estriol-3-glucuronide may optionally be used as the first analogue*".

Thus, whilst WO 96/16322 refers to the use of peptide epitope mimics of target ligands, and also refers to steroid target ligands, there is no explicit disclosure or suggestion of a peptide mimic for a steroid target ligand. Indeed the only mention of particular analogues for steroid target ligands are other, closely-related, steroids. Accordingly, the person skilled in the art would not deduce from the content of WO 96/16322 that peptide mimics for steroid analogues actually existed or could be made, and would not have any reasonable expectation of success in this regard as there is no evidence to suggest that such existed.

A brief reference, along similar lines to that discussed above, appears in WO 99/27356 (page 10) but again without specific examples or any experimental evidence.

Saviranta *et al.*, (1998 Bioconjugate Chem. 9, 725-735) disclose an assay for estradiol using Fab fragments specific for estradiol. There is no mention or suggestion of a peptide mimotope of estradiol which binds to the Fab fragments.

Slootstra *et al.*, (1997 *Journal of Molecular Recognition* 10, 217-224) describe a screening method to identify synthetic peptides that mimic epitopes, but those authors only ever refer to mimics of protein or peptide antigens, and there is no recognition or suggestion that peptide mimotopes might be available in respect of steroid compounds.

Lastly, US 5,635,182 (McCoy & Lu) relates to subject matter very different to the present invention and is generally concerned with DNA sequences encoding thioredoxin-like polypeptides. There is a brief disclosure (Seq. ID No. 2 therein) of a 20mer peptide which includes the tripeptide sequence Phe-Glu-Asp.

Summary of the Invention

This invention is based on the unexpected discovery that despite significant structural differences, peptide mimotopes for certain steroidal compounds do in fact exist and can be advantageously used, for example, in competitive or displacement-type immunoassays designed for the detection of steroids in a sample. In this regard, the present invention is directed to a purified peptide mimotope which is capable of binding specifically to an antibody specific to estradiol, and to isolated nucleic acid sequences encoding the purified peptide mimotope. It is also directed to an immunoassay test device for the detection in a sample of estradiol, the immunoassay comprising the peptide mimotope, as well as an antibody capable of binding specifically to the peptide mimotope to generate a detectable signal.

The present invention provides numerous advantages. In addition to the peptides being capable of being utilised as immunogens, the peptide mimotopes can be used to construct new, or improve the performance of old, immunoassay test formats and devices. They can, for example, be utilised essentially to "tune" the signal in conventional displacement assays for the detection of estradiol. Further, they can be bound directly to certain assay surfaces which are otherwise non-compatible with estradiol, the estradiol on such surfaces needing to be bound to the surface by complexing with another -- often proteinaceous -- molecule. Other advantages will become readily apparent in the description of the invention below.

Detailed Description of the Invention

The peptide mimotopes of the invention are capable of specific binding to any antibody which is specific to estradiol. Estradiol as used herein shall be taken to mean estradiol or metabolites thereof (e.g. the preferred estrone-3-glucuronide), as well as any related steroidal compounds having a basic estrone structure. Such related compounds are exemplified by, but not necessarily limited to, estriol, 16-epiestriol, 17-epiestriol, 17- β -estradiol 3-(β -D-glucuronide), estriol 3-(β -D-glucuronide), estrone, 17 α -ethynylestradiol, and 16 α -hydroxyestrone.

By specific binding it is meant that the mimotope is capable of being bound to the antigen-binding site of an antibody in a selective fashion in the presence of excess quantities of other materials not of interest, and tightly enough (i.e. with high enough affinity) that when used in an immunoassay, it provides a useful assay result. Similarly, an antibody "specific to estradiol" is one which is capable of binding to estradiol (or related compounds) in a selective fashion in the presence of excess quantities of other materials not of interest, and tightly enough that when used in an immunoassay it provides a useful assay result.

The antibody to which the peptide mimotopes are capable of being specifically bound can be any antibody, fragment or construct thereof, having a binding specificity for estradiol or metabolites thereof. Various forms of such antibodies are contemplated which may include monoclonal or polyclonal antibodies, Fv, Fab, ScFv and the like. Also contemplated are multivalent and/or multispecific constructions which have been described in the literature and comprise two or more polypeptide chains -- see for example, patent application Harris et al., WO 94/09131 and Davis et al., WO 97/14719 -- or are based on a 'double ScFv' approach, wherein the multivalency arises when two or more monovalent ScFv molecules are linked together, providing a single chain molecule comprising at least four variable domains, as described, for example, in Whitlow et al., WO 93/11161 and Mezes et al., WO 94/13806.

The antibodies, when utilized with the peptide mimotopes in an immunoassay test device, can be constructed by methods known in the art. Techniques such as those exemplified in Verhoeven and Windust, *Advances in Antibody Engineering in Molecular Immunology: Frontiers in Molecular Biology*, 2nd Ed., published by Oxford University Press, pp. 283-325 (Oxford, 1995) and Price et al. *Principles and Practice of Immunoassays*, 2nd Ed., published by Macmillan Publishers Ltd (London, 1997) are suitable. Many antibodies may also be obtained commercially. For the estradiol metabolite, estrone-3-glucuronide, a monoclonal antibody is described in Linscott's *Directory of Immunological and Biological Reagents* (10th edition 1998-9) and may be obtained from OEM Concepts Inc, Toms River, NJ, USA.

As described in the accompanying examples, the peptide mimotopes of the invention have been identified from epitope libraries by various screening techniques. They have also been identified from peptide libraries constructed from the known naturally occurring amino acids.

The peptide mimotopes will contain a minimum core binding region. That is, they will include a minimum continuous amino acid sequence which is necessary for imparting to the mimotope the capability of specific binding to the target antibody. Preferably, the region is such that the affinity of the mimotope for the binding reaction to a single antibody binding site in solution is greater than or equal to 10^5 L/mole. Methods of determining binding affinity are routine for those skilled in the art, and the binding affinity of a particular mimotope for a particular antibody can be readily measured with the benefit of the techniques described herein and using the common general knowledge of those skilled in the art.

The peptide mimotopes can be any size, though it is preferable that they be smaller than that which would allow for tertiary or globular structuring to occur. Thus, they are typically no larger than 30, and preferably no greater than 20, amino acids in length. The core binding region of each mimotope will typically be less than 12 amino acids, preferably less than 7 amino acids, and optimally between 3 to 6 amino acids in length. Preferred mimotopes are identified in the examples below.

In particular, the inventors have found that the core binding region of many mimotopes in accordance with the invention comprises one of the three tripeptide sequences identified as follows: Xaa-Glu-Asp; Phe-Xaa-Asp; and Phe-Glu-Xaa. Thus, in general, preferred mimotopes will comprise one of these three tripeptide sequences (typically, Phe-Glu-Asp), but it should be noted that mere possession of such a tripeptide is not necessarily sufficient for the peptide to possess suitable specific binding activity: the inventors have found some examples of peptides which comprise one of the aforementioned tripeptide sequences but which do not exhibit suitable specific binding activity. With the benefit of the present disclosure, those skilled in the art will readily be able to screen candidate peptides and select those having the most desirable binding characteristics.

It will be apparent from the present disclosure that, if using D-isomers of amino acids, the reverse sequences may be employed (i.e. Xaa-Glu-Phe; Asp-Xaa-Phe; and Asp-Glu-Xaa).

US 5,635,182 discloses a 20mer peptide derived from bovine phospholipase C-II, having the sequence QPFEDFRISQEHLADHFDGR.

The present inventors have found that several peptides comprising the tripeptide FED are useful as peptide mimotopes in accordance with the invention. The inventors have not performed any experiments to investigate whether the 20mer disclosed in US 5,635,182 might also be useful (i.e. be capable of binding specifically to an antibody specific to estradiol) but, in the event that the prior art peptide does exhibit such specific binding activity, the inventors hereby provisionally disclaim the peptide consisting of the amino acid sequence QPFEDFRISQEHLADHFDGR.

Purification of the peptide mimotopes can be accomplished by conventional means, such as those described in Tendler et al., *The role of the arginine residue in the stabilization of mucin core type I β turns*. Protein and Peptide Letters, 1, 39-43 (1994). Preferably, the peptide mimotopes will be purified to 95%, optimally to 99%. The mimotope is typically provided as a simple peptide, but may optionally be covalently peptide bonded or linked in some other way to other moieties, such as a label or a solid support.

In one embodiment of the invention, the peptide mimotopes are utilized in an immunoassay test device. Such a device can take different forms, and it can be varied depending on the precise nature of the assay being performed.

Because the peptide mimotopes "mimic" a substance (i.e. Estradiol) which will often be the subject of testing, they are essentially antigenic by nature and function. Thus, it is most preferable that they be utilized in competitive or displacement-type assays (hereinafter collectively referred to as competitive assays). Nothing, however, would preclude their usage in conventional sandwich-type assays as well and specific formats can be readily designed.

Specifically, it is contemplated that in a competitive assay incorporating the peptide mimotopes of the invention, the mimotopes would be coated onto a solid support, typically nitrocellulose or other hydrophobic porous material. They may also be coated on synthetic plastics materials, microtitre assay plates, latex beads, filters comprising cellulosic or synthetic polymeric materials, glass or plastic slides, dipsticks, capillary fill devices and the like.

Coating of the peptide mimotopes to these surfaces can be accomplished by methods known in the art and described in, for example, EP-B-0291194. A particular advantage of the present invention is that unlike the compounds which they mimic, the mimotopes of invention are peptides, and thus can be coated directly onto certain assay surfaces such as nitrocellulose. Estradiol, by contrast, is non-compatible with such cellulosic materials and thus often needs to be bound to the surface by forming a complex with another molecule. Proteins are typically used for such complexing, with BSA often being the most preferred.

In a preferred competitive assay the peptide mimotopes, once coated on the surface of a support, are specifically bound to antibodies or fragments or constructs thereof. The antibodies can be as described above and should be capable of specific binding to estradiol. It is envisioned that a liquid sample containing estradiol migrating over the region

containing the antibodies bound to the mimotopes would displace a certain amount of antibodies from the surface of the support. The amount of antibodies displaced would be dependent on several factors including the concentration of the estradiol in the sample, and the relative binding affinities of the mimotopes and estradiol for the antibodies. The amount of antibody displaced could then be measured as a means to determine the relative concentration of estradiol in the sample.

Alternatively and in another preferred embodiment, it is contemplated that the antibodies could be bound to the surface, with the peptide mimotopes being specifically bound to the antibodies and capable of being displaced by estradiol migrating in a sample in contact with (e.g. through) the support. The displacement would generate a measurable signal of the amount of peptide mimotopes displaced and hence the amount of estradiol in the sample.

Other immunoassay test devices contemplated by the invention include those employing, for example, capillary-fill means in which a liquid sample is drawn into a device by capillary action along a suitably-proportioned capillary inlet. Capillary-fill devices which may be adapted for use in the present invention are disclosed, for example, in Shanks et al., U.S. Patent 5,141,868, Shanks et al., EP-A-0422708, and Birch et al., EP-B-0274215.

Devices such as those described in May et al., U.S. Patent 5,622,871 and May et al., U.S. Patent 5,656,503 are also suitable for practice of the immunoassays of the invention. If used, these devices preferably comprise a hollow elongated casing containing the solid support. The solid support communicates indirectly with the exterior of the casing via a bibulous fluid sample receiving member which may or may not protrude from the casing, the solid support and the sample receiving member being linked so as to allow for the fluid sample to migrate between the two by capillary action.

Spatially distant along the solid support from the sample receiving member are the test and, optionally, control zones. Within the test zone, the peptide mimotopes can be bound to an antibody immobilized on the support. Such immobilisation can be accomplished by

any number of known means including chemically coupling using, for example, CNBr, carbonyldiimidazole, or tresyl chloride. Alternatively, various "printing" techniques may be used. These include application of liquid antibodies by micro-syringes, direct printing, ink-jet printing, and the like. Chemical or physical treatment of the support prior to application of the antibody is also specifically contemplated, as such may facilitate immobilisation.

The casing in such devices is typically constructed of opaque or translucent material incorporating at least one aperture through which the analytical result may be observed, either by the naked eye or electronic means.

Such devices can be provided to clinical laboratories or as kits suitable for home use, such kits comprising one or more devices individually wrapped in moisture impervious wrapping and packaged together with appropriate instructions to the user.

The sample receiving member can be made from any bibulous, porous or fibrous material capable of absorbing liquid rapidly. The porosity of the material can be unidirectional (i.e. with pores or fibres running wholly or predominantly parallel to an axis of the member) or multidirectional (omnidirectional, so that the member has an amorphous sponge-like structure). Porous plastics material, such as polypropylene, polyethylene (preferably of very high molecular weight), polyvinylidene fluoride, ethylene vinylacetate, acrylonitrile and polytetrafluoro-ethylene can be used. It can be advantageous to pre-treat the member with a surface-active agent during manufacture, as this can reduce any inherent hydrophobicity in the member and therefore enhance its ability to take up and deliver a moist sample rapidly and efficiently. Porous sample receiving members can also be made from paper or other cellulosic materials, such as nitrocellulose. Preferably the material comprising the sample receiving member should be chosen such that the porous member can be saturated with liquid sample within a matter of seconds. The liquid must be capable of permeating freely from the porous sample receiving member into the solid support.

The solid support in such devices is preferably a dry porous carrier. It may be made of separate strips or sheets and, like the sample receiving member, can be constructed from any material capable of allowing the liquid sample to migrate through a portion of its length by, preferably, capillary action. The support should allow for the immobilisation of the antibody and/or peptide mimotope on its surface, and should not interfere with the binding reactions which are necessary for the proper functioning of the assay.

The solid support may have associated with it an absorbent "sink" which will facilitate capillary action of fluid up the length of the support, and will provide a means by which to avoid flooding of the test device by application of excess sample. Specific materials for and applications of sinks are conventional in the art and may be readily applied to the devices of the present invention.

In the immunoassay test devices of the invention, in order to provide a measurable signal of the amount of analyte in the sample it is preferred that either the peptide mimotope, or the antibody to which it is bound, be labelled. In the preferred embodiment of the invention, the label is any entity the presence of which can be readily detected. Preferably the label is a direct label, such as the those described in detail in May et al., U.S. Patent 5,656,503. Direct labels are entities which, in their natural state, are readily visible either to the naked eye, or with the aid of an optical filter and/or applied stimulation, e.g. UV light to promote fluorescence. Examples include radioactive, chemiluminescent, electroactive (such as redox labels), and fluorescent compounds. Direct particulate labels, such as dye sols, metallic sols (e.g. gold) and coloured latex particles, are also very suitable and are, along with fluorescent compounds, preferred. Of these options, coloured latex particles and fluorescent compounds are most preferred. Concentration of the label into a small zone or volume should give rise to a readily detectable signal, e.g. a strongly coloured area.

Indirect labels, such as enzymes, e.g. alkaline phosphatase and horseradish peroxidase, can also be used, but these usually require the addition of one or more developing reagents such as substrates before a visible signal can be detected. Hence, they are less preferred.

Such additional reagents can be incorporated in the solid support of the assay device such that they dissolve or disperse when a liquid sample is applied. Alternatively, the developing reagents can be added to the sample before application of the sample to the solid support.

Conjugation of the label to the peptide mimotope or the antibody can be by covalent or non-covalent (including hydrophobic) bonding, or by adsorption. Techniques for such conjugation are commonplace in the art and may be readily adapted for the particular reagents employed. In the preferred embodiment wherein the label is a coloured latex particle, the label is preferably conjugated to the antibody and it is accomplished through adsorption. Where the label is a fluorescent compound, it is preferred that the label be conjugated to or constructed as part of the antibody.

Upon usage of the test device, the label can provide a test and/or control signal which can be detected from the test and control surfaces by known conventional means. This includes evaluation by the naked eye, or more typically when precise measurements are desired, by appropriate instrumentation. Instrumentation is particularly suitable when the control or test signal is measured by the amount of mass of complex at the control or test surface.

The immunoassay test devices of the invention may be applied to virtually any type of biological or non-biological sample, though liquid biological samples derived from urine or serum are preferred. The samples may be purified or diluted prior to assaying. The term "immunoassay test device" as used herein is also intended to encompass components of immunoassay test devices which may be sold or supplied as separate articles and which require the presence of other components in order to form a working test device. In particular, it is contemplated that the immunoassay test device of the invention may be a dipstick, test stick or the like, which are generally provided as disposable items and may be supplied as separate components.

In a second aspect the invention provides an isolated nucleic acid encoding a peptide mimotope in accordance with the first aspect of the invention. The nucleic acid may be

prepared by cloning from a library of sequences or from an organism (e.g. a phage or a bacterium), or prepared by *in vitro* synthesis using standard techniques (e.g. automated solid phase oligonucleotide synthesisers, which are commercially available from many sources) or, less conveniently, by performance of ligation reactions, ligating together component nucleic acid sequences from different sources. Typically the isolated nucleic acid sequence will be a DNA sequence (but could, conceivably, be a sense RNA sequence) and will comprise a minimum of 9 bases. More typically, the nucleic acid (or rather, that portion thereof which encodes the peptide mimotopes) will comprise between 12 and 90 bases, desirably between 15 and 90, and preferably between 15 and 60 bases. The nucleic acid may advantageously comprise other components, such as promoter, enhancer and terminator sequences, one or more origins of replication, and the like. In addition, the isolated nucleic acid may encode a fusion protein, in which the peptide mimotope is fused (at either the 5' or 3' terminus) to another polypeptide moiety such as a polypeptide label. In such embodiments as aforesaid, whilst that portion of the nucleic acid which encodes the mimotope will generally comprise a number of bases within the ranges identified above, the nucleic acid as a whole may be considerably larger. It will be understood that the nucleic acid molecule may, in some embodiments, encode a peptide which consists solely of the peptide mimotope without any extraneous amino acid residues (e.g. the mimotope will be in isolation from the sequences adjacent thereto in any naturally-occurring molecule from which the mimotope may be derived).

The isolated nucleic acid molecule may conveniently take the form of a plasmid or other replicable moiety.

In a third aspect, the invention provides for the use of a peptide mimotope in accordance with the first aspect of the invention defined above, to assay for the presence and/or amount of estradiol in a sample to be tested.

The practice of the invention is described in detail below with reference to specific illustrative examples, but the invention is not to be construed as being limited thereto.

EXAMPLES

Identification of Peptide Mimotopes for Estradiol

Means by which to identify examples of peptide mimotopes of the estradiol metabolite, estrone-3-glucuronide, are described below.

Monoclonal antibodies MAb 4155 were expressed from the 4155 monoclonal cell line. The 4155 monoclonal cell line was prepared and screened according to the methods described by Gani et al., (J Steroid Biochem. Molec. Biol. 48, 277-282 (1994)). The Gani et al. publication relates to development of anti-progesterone antibodies, but similar techniques were employed in producing antibodies reacting with estrone and analogues thereof.

Comparative amino acid sequences utilized in the following examples are as follows:

| | |
|-------------------------------------|----------------|
| Glu-Asp | (SEQ ID NO:5) |
| Ala-Ala-Glu-Arg-Gly-Leu-Phe-Glu | (SEQ ID NO:71) |
| Ala-Ala-Glu-Arg-Gly-Leu-Phe | (SEQ ID NO:72) |
| Ala-Ala-Glu-Arg-Gly-Leu | (SEQ ID NO:73) |
| Ala-Ala-Glu-Arg-Gly | (SEQ ID NO:74) |
| Ala-Ala-Glu-Arg | (SEQ ID NO:75) |
| Ala-Ala-Glu | (SEQ ID NO:76) |
| Ala-Ala | (SEQ ID NO:77) |
| Ala-Ala-Glu-Arg-Gly-Leu-Ala-Glu-Asp | (SEQ ID NO:78) |
| Ala-Ala-Glu-Arg-Gly-Leu-Phe-Ala-Asp | (SEQ ID NO:79) |
| Ala-Ala-Glu-Arg-Gly-Leu-Phe-Glu-Ala | (SEQ ID NO:80) |

EXAMPLE 1

Identification of Peptide Mimotope Sequences by Phage Display

pVIII9aa-cys nonapeptide phage library

The VIII9aa-cys library phage library described by Felici F et al., *Mimicking of discontinuous epitopes by phage-displayed peptides, II. Selection of clones recognised by a protective monoclonal antibody against the Bordetella pertussis toxin from phage peptide libraries*. Gene 128, 21-27 (1993) and Luzzago et al. *Mimicking of discontinuous epitopes by phage-displayed peptides, I. Epitope mapping of human H ferritin using a phage library of constrained peptides*. Gene 128, 51-57 (1993) was used. The library consisted of random nonapeptides fused to the major coat protein pVIII so that several hundred peptides were displayed on each phage particle.

Screening of phage library

Affinity selection of phage was performed by a combination of the methods of Folgori A. et al. *A general strategy to identify mimotopes of pathological antigens using only random peptide libraries and human sera*. EMBO J 13, 2236-2243 (1994) and Parmley S.F. et al. *Antibody-selectable fd phage vectors: affinity purification of target genes*. Gene 73, 305-318 (1988).

Polystyrene tubes used for panning (Immunotubes™ from Nunc) were coated either with affinity-purified anti-estrone-3-glucuronide antibodies (20 µg) in 2 mls of coating buffer (0.1 M NaHCO₃, pH 9.0) or with coating buffer only overnight at 4° C. After three washes with tris buffered saline (TBS; 50 mM tris-HCl, 140 mM NaCl, pH 7.4) both tubes were incubated with 4 mls of blocking buffer (TBS containing 10 mg/ml ovalbumin) for 4 h at room temperature. The VIII9aa-cys library was shown to have a titre of 1×10^{13} transducing units/ml (TU/ml) by infection of logarithmic XL1-Blue bacteria (Stratagene, Amsterdam, Holland). Aliquots (1 µl; 1×10^{11} TU) from the donated phage suspension were added to the antibody-coated and un-coated polystyrene tubes each containing 1 ml of TBS and 1mg/ml ovalbumin and incubated overnight at 4°C. Unbound phage were removed by 15 washes (each of 4 ml) with TBS containing 0.5% (v/v) Tween 20™ (TTBS) followed by 5 washes with TBS at room temperature. Bound phage were eluted by incubation of washed panning tubes with 1ml of elution buffer (0.1 M HCl, pH 2.2,

adjusted with glycine, containing 1 mg/ml ovalbumin) for 12 min at room temperature. The eluted phage were transferred to 1 ml polypropylene tubes and neutralised with 60 μ l of 2 M tris (pH not adjusted). Aliquots (200 μ l) of 1 M tris-HCl, pH 7.4 were also added to the panning tubes for neutralisation. The eluted neutralised phage particles (1 ml) were used for infection of 9 ml of logarithmic XL1-Blue bacteria (in 2TY containing 1% (w/v) glucose). Logarithmic XL1-Blue bacteria (4 ml) were also added directly to the neutralised panning tubes. Infection was carried out for 30 min at 37°C with no shaking. The infected bacteria were then pooled (total volume 13 ml) ampicillin was added (to 100 μ g/ml) and the cultures incubated overnight with shaking at 37°C. A small aliquot (10 μ l) of infected bacterial cells was removed prior to overnight incubation for titration (diluted 10^{-2} to 10^{-6} in 2TY/Amp/Glucose) on 2TY agar containing 1% (w/v) glucose and ampicillin (100 μ g/ml). An aliquot (150 μ l) of the overnight XL1- Blue culture infected with phage eluted from the panning tube coated with MAb 4155 antibodies was then added to 15 ml of 2TY containing 1% (w/v) glucose and 100 μ g/ml ampicillin and grown to logarithmic phase. The cells were then superinfected with M13K07 helper phage (Gibco BRL Life Technologies, Paisley, Scotland) (1×10^{11} phage/ml) and incubated for 30 min without shaking at 37°C. This was followed by centrifugation for 20 min at 1800 rpm and resuspension of the cell pellet in 200 ml of 2TY containing 100 μ g/ml ampicillin and 25 μ g/ml kanamycin. The bacterial culture was then incubated overnight at 37° C with shaking. Bacterial cells were then pelleted by centrifugation (5000 rpm; 15 min) and the phage particles in the supernatant precipitated by addition of 40 mls of PEG/NaCl (2.5M NaCl containing 20% (w/v) polyethylene glycol 8000) and incubation on ice for 1 hour. The phage suspension was then spun at 10,000rpm for 20 min at 4°C and the resulting pellet resuspended in 20 ml of TBS. A further PEG precipitation was carried out by addition of 4 ml of PEG/NaCl and incubation on ice for a further 20 min. The final phage pellet was dissolved in 2 ml of TBS which resulted in phage titres of the order of 1×10^{13} TU/ml. These phage particles were added directly to fresh panning tubes and the entire panning procedure repeated a further two times. The entire screening protocol (three rounds of panning) was repeated after the first screen.

Phage ELISAs

The output from the third round of panning was plated out on 2TY agar, ampicillin (100 $\mu\text{g/ml}$) and 1% (w/v) glucose and incubated overnight at 37°C. Random individual bacterial colonies (~200) were picked and added to the wells of 96-well microtitre plates (Sterilin™) each containing 200 μl of 2TY, 1% (w/v) glucose and ampicillin (100 $\mu\text{g/ml}$). The microtitre plates were incubated overnight with shaking at 37°C. The following day aliquots from each well (20 μl) were added to the wells of fresh microtitre plates each containing 200 μl of 2TY, 1% glucose, 100 $\mu\text{g/ml}$ ampicillin and incubated with shaking for 1 h at 37°C. At the next stage, 25 μl of 2TY containing ampicillin (100 $\mu\text{g/ml}$), 1% (w/v) glucose and 10^9 M13KO7 helper phage were added to each well. The plates were incubated for 30 min at 37°C without shaking followed by a further incubation for 1 h at 37°C with shaking. The plates were then spun at 1800 rpm for 20 min at room temperature, the supernatant aspirated off and the cell pellet resuspended in 200 μl of 2TY containing ampicillin (100 $\mu\text{g/ml}$) and kanamycin (20 $\mu\text{g/ml}$). Incubation with shaking at 37°C was then carried out overnight. Centrifugation of overnight cultures in the wells of microtitre plates was carried out (1500 rpm; 20 mins) and phage-containing supernatants (100 μl) were added to sheep anti-M13 bacteriophage (C.P. Laboratories, Bishops Stortford, UK) coated microtitre plates (Greiner™, high bind). Purified sheep anti-M13 antibody-coated plates were prepared by overnight incubation (100 $\mu\text{l/well}$; 10 $\mu\text{g/ml}$) at 4°C in binding buffer (0.1 M NaHCO₃, pH 9.0). Blocking was carried out with PBST containing 10 mg/ml ovalbumin (200 $\mu\text{l/well}$) for 1 h at room temperature. After removal of unbound phage from sheep anti-M13-coated plates by five washes with PBST affinity-purified anti-estrone-3-glucuronide antibodies were added (20 μg) in 2 mls of PBST containing 10 mg/ml ovalbumin; 100 μl per well). Incubation was carried out for 2 h at room temperature. Alkaline phosphatase conjugated rabbit anti-mouse immunoglobulin (100 $\mu\text{l/well}$) was then added at a dilution of 1/1000 (in PBST, 10 mg/ml ovalbumin) and incubated for a further 2 h at room temperature. The assay was developed with 100 $\mu\text{l/well}$ of *p*-nitrophenyl phosphate (1 mg/ml) in 1M diethanolamine, 1 mM MgCl₂, pH 9.6 and the plates read at 410 nm.

DNA Sequencing

Double-stranded phagemid DNA was purified from bacterial cultures (50 ml) infected with positive phage clones using the Qiagen™ plasmid purification kit according to the manufacturer's instructions. Sequencing was carried out on an Applied Biosystems automated sequencer (Model 373A, version 1.2.0) using the oligonucleotide primer SEQ ID NO 1:

5'- TTT CCC AGT CAC GAC GTT G -3' (SEQ ID NO:1).

From the Phage ELISA and DNA sequencing results the following three peptide mimotope sequences were identified:

Ala-Ala-Glu-Arg-Gly-Leu-Phe-Glu-Asp (SEQ ID NO:2).

Thr-Ala-Trp-Thr-Tyr-Val-Leu-Gly-Phe (SEQ ID NO:3).

Thr-Ser-Trp-Ala-Tyr-Val-Leu-Gly-Pro (SEQ ID NO:4).

Identification of Mimotope Core Binding Regions by Replacement Net Analysis

Solid Phase Peptide Synthesis on Pins

Peptides were synthesised in duplicate or triplicate from the C-terminus by solid phase peptide synthesis on the heads of polyethylene pins (Geysen et al., *Strategies for epitope analysis using peptide synthesis*. J. Immunol. Methods 102, 259-274 (1987)) using a Multipin Peptide Synthesis Kit (Chiron Mimotopes, Victoria, Australia). Pins were arranged in a plastic holder in the format of a 96-well microtitre plate.

ELISA Testing of Peptides on Pins

All incubation steps were performed at room temperature (18-25°C) by lowering the pins reagents dispensed into 96-well microtitre plates (Becton Dickinson, CA, USA). Washing

was accomplished by placing the block of pins in a bath of phosphate buffered saline (PBS) containing Tween 20™ (0.01% v/v) with agitation for four cycles of 5 min. Non-specific binding sites on the surface of the pins were blocked by incubating in PBS containing casein (1% w/v, 175 µl/well) for 1 h. MAb4155 was diluted in blocking buffer and the pins were incubated in the antibody solution (150 µl/well) for 18 h at 4°C. After washing, the pins were incubated in horseradish peroxidase (HRP)-conjugated rabbit anti-mouse immunoglobulin (Dako, High Wycombe, UK, 1/1000 in blocking buffer for 1 h at 150 µl/well). The pins were washed once more and then incubated in ABTS [2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid)] working substrate for 15 min (150 µl/well). ABTS was prepared as a 0.033% (w/v) solution in 0.1M citrate phosphate buffer (pH 4.5) with 33% hydrogen peroxide (1 µl/ml). Colour development was terminated by removal of the pins from the wells and was measured spectrophotometrically at 405 nm using a Milenia Kinetic Analyser™ (DPC, Llanberis, Wales).

Identification of the Core Binding Regions

In order to identify core binding regions, a set of peptides was synthesised on the heads of pins based on SEQ ID NO:2. These included peptides sequentially reduced in length by one amino acid, first from the N-terminus and then, in another series, from the C-terminus. In addition, a set of peptides was synthesised in which each residue of the lead sequence was replaced by Ala (or Gly if Ala already existed at that position) in order to assess the contribution of each residue to the binding event. MAb4155 was tested for reactivity with these peptides, the results being shown below in Table 1.

Table 1

| Mimotope | SEQ ID NO: | Type | Relative Binding* |
|-------------------------------------|------------|-----------|-------------------|
| Ala-Ala-Glu-Arg-Gly-Leu-Phe-Glu-Asp | 2 | invention | 2.0 |
| Ala-Glu-Arg-Gly-Leu-Phe-Glu-Asp | 11 | invention | 1.5 |
| Glu-Arg-Gly-Leu-Phe-Glu-Asp | 10 | invention | 1.8 |
| Arg-Gly-Leu-Phe-Glu-Asp | 9 | invention | 1.6 |

| | | | |
|-------------------------------------|----|------------|-----|
| Gly-Leu-Phe-Glu-Asp | 8 | invention | 2.2 |
| Leu-Phe-Glu-Asp | 7 | invention | 1.9 |
| Phe-Glu-Asp | 6 | invention | 2.0 |
| Glu-Asp | 5 | comparison | 1.0 |
| Ala-Ala-Glu-Arg-Gly-Leu-Phe-Glu | 71 | comparison | 1.0 |
| Ala-Ala-Glu-Arg-Gly-Leu-Phe | 72 | comparison | 1.0 |
| Ala-Ala-Glu-Arg-Gly-Leu | 73 | comparison | 1.0 |
| Ala-Ala-Glu-Arg-Gly | 74 | comparison | 1.0 |
| Ala-Ala-Glu-Arg | 75 | comparison | 0.8 |
| Ala-Ala-Glu | 76 | comparison | 0.9 |
| Ala-Ala | 77 | comparison | 1.0 |
| Gly-Ala-Glu-Arg-Gly-Leu-Phe-Glu-Asp | 12 | invention | 1.9 |
| Ala-Gly-Glu-Arg-Gly-Leu-Phe-Glu-Asp | 13 | invention | 2.0 |
| Ala-Ala-Ala-Arg-Gly-Leu-Phe-Glu-Asp | 14 | invention | 1.5 |
| Ala-Ala-Glu-Ala-Gly-Leu-Phe-Glu-Asp | 15 | invention | 2.0 |
| Ala-Ala-Glu-Arg-Ala-Leu-Phe-Glu-Asp | 16 | invention | 1.5 |
| Ala-Ala-Glu-Arg-Gly-Ala-Phe-Glu-Asp | 17 | invention | 1.6 |
| Ala-Ala-Glu-Arg-Gly-Leu-Ala-Glu-Asp | 78 | comparison | 0.8 |
| Ala-Ala-Glu-Arg-Gly-Leu-Phe-Ala-Asp | 79 | comparison | 1.8 |
| Ala-Ala-Glu-Arg-Gly-Leu-Phe-Glu-Ala | 80 | comparison | 1.1 |

* Relative Binding of MA4155 to peptides as measured by ELISA as described above

As is demonstrated from the data, of the peptides tested, only those amino acid sequences comprising the core binding region as indicated by SEQ ID NO:6 provided adequate binding. Amino acid sequences represented by SEQ ID NO:7 - 17 contain the core binding region of SEQ ID NO:6 and provided adequate binding to serve as an estradiol mimotope.

Phe-Glu-Asp

(SEQ ID NO:6)

Leu-Phe-Glu-Asp

(SEQ ID NO:7)

| | |
|-------------------------------------|----------------|
| Gly-Leu-Phe-Glu-Asp | (SEQ ID NO:8) |
| Arg-Gly-Leu-Phe-Glu-Asp | (SEQ ID NO:9) |
| Glu-Arg-Gly-Leu-Phe-Glu-Asp | (SEQ ID NO:10) |
| Ala-Glu-Arg-Gly-Leu-Phe-Glu-Asp | (SEQ ID NO:11) |
| Gly-Ala-Glu-Arg-Gly-Leu-Phe-Glu-Asp | (SEQ ID NO:12) |
| Ala-Gly-Glu-Arg-Gly-Leu-Phe-Glu-Asp | (SEQ ID NO:13) |
| Ala-Ala-Ala-Arg-Gly-Leu-Phe-Glu-Asp | (SEQ ID NO:14) |
| Ala-Ala-Glu-Ala-Gly-Leu-Phe-Glu-Asp | (SEQ ID NO:15) |
| Ala-Ala-Glu-Arg-Ala-Leu-Phe-Glu-Asp | (SEQ ID NO:16) |
| Ala-Ala-Glu-Arg-Gly-Ala-Phe-Glu-Asp | (SEQ ID NO:17) |

Additional core binding sequences were identified utilizing the amino acid sequence SEQ ID NO:8 and investigating the effect of systematic replacement of each residue by the other 19 naturally occurring amino acids using known techniques as exemplified in Verhoeven et al., *Construction of a reshaped HMFG1 antibody and comparison of its fine specificity with that of the parent mouse antibody*. Immunology, 78, 364-370 (1993).

The sequences which had superior binding reactivity and specificity compared to SEQ ID NO:8 are identified as follows. The binding of MAb4155 to these sequences as determined by ELISA is shown below in Table 2, setting SEQ ID NO:8 to a Relative Binding of 100.

| | |
|---------------------|----------------|
| Gly-Phe-Phe-Glu-Asp | (SEQ ID NO:18) |
| Gly-Trp-Phe-Glu-Asp | (SEQ ID NO:19) |
| Gly-Tyr-Phe-Glu-Asp | (SEQ ID NO:20) |
| Gly-Leu-Trp-Glu-Asp | (SEQ ID NO:21) |
| Gly-Leu-Phe-Cys-Asp | (SEQ ID NO:22) |
| Gly-Leu-Phe-Asp-Asp | (SEQ ID NO:23) |
| Gly-Leu-Phe-Phe-Asp | (SEQ ID NO:24) |
| Gly-Leu-Phe-Ile-Asp | (SEQ ID NO:25) |
| Gly-Leu-Phe-Leu-Asp | (SEQ ID NO:26) |
| Gly-Leu-Phe-Trp-Asp | (SEQ ID NO:27) |

| | |
|---------------------|----------------|
| Gly-Leu-Phe-Tyr-Asp | (SEQ ID NO:28) |
| Gly-Leu-Phe-Glu-Cys | (SEQ ID NO:29) |
| Gly-Leu-Phe-Glu-Phe | (SEQ ID NO:30) |
| Gly-Leu-Phe-Glu-Ile | (SEQ ID NO:31) |
| Gly-Leu-Phe-Glu-Leu | (SEQ ID NO:32) |
| Gly-Leu-Phe-Glu-Val | (SEQ ID NO:33) |
| Gly-Leu-Phe-Glu-Trp | (SEQ ID NO:34) |
| Gly-Leu-Phe-Glu-Tyr | (SEQ ID NO:35) |

100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2238
2239
2240
2241
2242
2243
2244
2245
2246
2247
2248
2249
2250
2251
2252
2253
2254
2255
2256
2257
2258
2259

Table 2

| Mimotope | SEQ ID NO: | Relative Binding* |
|---------------------|------------|-------------------|
| Gly-Leu-Phe-Glu-Asp | 8 | 100 |
| Gly-Phe-Phe-Glu-Asp | 18 | 200 |
| Gly-Trp-Phe-Glu-Asp | 19 | 343 |
| Gly-Tyr-Phe-Glu-Asp | 20 | 220 |
| Gly-Leu-Trp-Glu-Asp | 21 | 207 |
| Gly-Leu-Phe-Cys-Asp | 22 | 335 |
| Gly-Leu-Phe-Asp-Asp | 23 | 121 |
| Gly-Leu-Phe-Phe-Asp | 24 | 184 |
| Gly-Leu-Phe-Ile-Asp | 25 | 169 |
| Gly-Leu-Phe-Leu-Asp | 26 | 138 |
| Gly-Leu-Phe-Trp-Asp | 27 | 578 |
| Gly-Leu-Phe-Tyr-Asp | 28 | 252 |
| Gly-Leu-Phe-Glu-Cys | 29 | 296 |
| Gly-Leu-Phe-Glu-Phe | 30 | 204 |
| Gly-Leu-Phe-Glu-Ile | 31 | 174 |
| Gly-Leu-Phe-Glu-Leu | 32 | 168 |
| Gly-Leu-Phe-Glu-Val | 33 | 177 |
| Gly-Leu-Phe-Glu-Trp | 34 | 594 |
| Gly-Leu-Phe-Glu-Tyr | 35 | 386 |

* Relative Binding of MAb4155 to peptides as measured by ELISA as described above

D-isomers of the Reverse Sequence of the Core Binding Regions.

The foregoing SEQ ID NO.'s 1-35 are L-isomers. It was also demonstrated that the D-isomers of the reverse sequences of those core binding regions identified above similarly function as effective mimotopes. For example, an amino acid sequence as described by SEQ ID NO:37 was prepared by the peptide synthesis methods described above. Binding affinity relative to the parent sequence (SEQ ID NO:28) was measured by the described testing methods. The results show the reverse sequence to have an equivalent relative binding affinity compared to the parent sequence.

The following therefore identify core binding sequences of peptides capable of functioning as mimotopes for estradiol. Each sequence contains D-isomers of the amino acids in the reverse sequence of one of those described above.

NOTE: SEQ ID NO:s 36-56 are D-isomers

| | |
|---------------------|----------------|
| Asp-Glu-Phe | (SEQ ID NO:36) |
| Asp-Tyr-Phe-Leu-Gly | (SEQ ID NO:37) |
| Asp-Glu-Phe-Phe-Gly | (SEQ ID NO:38) |
| Asp-Glu-Phe-Trp-Gly | (SEQ ID NO:39) |
| Asp-Glu-Phe-Tyr-Gly | (SEQ ID NO:40) |
| Asp-Glu-Trp-Leu-Gly | (SEQ ID NO:41) |
| Asp-Cys-Phe-Leu-Gly | (SEQ ID NO:42) |
| Asp-Asp-Phe-Leu-Gly | (SEQ ID NO:43) |
| Asp-Phe-Phe-Leu-Gly | (SEQ ID NO:44) |
| Asp-Ile-Phe-Leu-Gly | (SEQ ID NO:45) |
| Asp-Leu-Phe-Leu-Gly | (SEQ ID NO:46) |
| Asp-Trp-Phe-Leu-Gly | (SEQ ID NO:47) |
| Cys-Glu-Phe-Leu-Gly | (SEQ ID NO:48) |
| Phe-Glu-Phe-Leu-Gly | (SEQ ID NO:49) |
| Ile-Glu-Phe-Leu-Gly | (SEQ ID NO:50) |
| Leu-Glu-Phe-Leu-Gly | (SEQ ID NO:51) |
| Val-Glu-Phe-Leu-Gly | (SEQ ID NO:52) |
| Trp-Glu-Phe-Leu-Gly | (SEQ ID NO:53) |

Tyr-Glu-Phe-Leu-Gly (SEQ ID NO:54)

Phe-Gly-Leu-Val-Tyr-Thr-Trp-Ala-Thr (SEQ ID NO:55)

Pro-Gly-Leu-Val-Tyr-Ala-Trp-Ser-Thr (SEQ ID NO:56)

EXAMPLE 2**Identification of Peptide Mimotope Sequences from Pepscan Libraries**

The pins from the Multipin Peptide Synthesis Kit as described in Example 1 were used to construct libraries of peptide sequences encompassing all possible trimer combinations of the 20 naturally occurring amino acids, supplemented by a further random set of dodecapeptides (Slootstra JW et al., *Screening of a small set of random peptides: a new strategy to identify synthetic peptides that mimic epitopes* J Molec. Recog. 10, 217-224 (1997)). The binding of MAb4155 was tested on the library for binding affinity in a manner as described in Example 1. This identified the following amino acid sequences (L-isomers) as core binding regions for estradiol mimotopes.

| | |
|---|----------------|
| Asp-Phe-Tyr | (SEQ ID NO:57) |
| Phe-Tyr-Glu | (SEQ ID NO:58) |
| Tyr-Glu-Glu | (SEQ ID NO:59) |
| Tyr-Gln-Glu | (SEQ ID NO:60) |
| Asn-Glu-Glu-Asp-Phe-Tyr-Gln-Ile-Gln-Leu-Tyr-Glu | (SEQ ID NO:61) |
| Arg-Gln-Ile-Asp-Phe-Tyr-Gln-Glu-Ile-Gln-Phe-Lys | (SEQ ID NO:62) |
| Asp-Asp-Phe-Tyr-Gly-Gln-Pro-Arg-Glu-Gln-Val-Arg | (SEQ ID NO:63) |

Similarly to Example 1 the following reverse sequences of D-amino acids are identified as capable of functioning as the core binding region for peptide mimotopes for estradiol.

NOTE: SEQ ID NO:s 64-70 are D-isomers

| | |
|---|----------------|
| Tyr-Phe-Asp | (SEQ ID NO:64) |
| Glu-Tyr-Phe | (SEQ ID NO:65) |
| Glu-Glu-Tyr | (SEQ ID NO:66) |
| Glu-Gln-Tyr | (SEQ ID NO:67) |
| Glu-Tyr-Leu-Gln-Ile-Gln-Tyr-Phe-Asp-Glu-Glu-Asn | (SEQ ID NO:68) |
| Lys-Phe-Gln-Ile-Glu-Gln-Tyr-Phe-Asp-Ile-Gln-Arg | (SEQ ID NO:69) |
| Arg-Val-Gln-Glu-Arg-Pro-Gln-Gly-Tyr-Phe-Asp-Asp | (SEQ ID NO:70) |

EXAMPLE 3

Comparison of Competitive Assays for Estrone-3-Glucuronide (E3G) Using E3G or a Peptide Mimotope of E3G on a Solid Phase

Peptide Ligand Synthesis

Synthetic peptide ligands were prepared on an Applied Biosystems 431A Peptide Synthesiser Biopolymer Synthesis and Analysis Unit, TM QMC, (Nottingham, UK). Purity was assessed by mass spectroscopy and HPLC and was in excess of 95%.

Preparation of Ovalbumin-E3G conjugate

An estrone-3-glucuronide (E3G) ovalbumin conjugate was prepared by resuspending 2.6mg of E3G in 2ml of freshly prepared solution of EDC (1-ethyl (dimethylaminopropyl) carbodiimide, 0.1M) and NHS (N-hydroxysuccinamide, 0.02M) and incubating for 15minutes at room temperature. To the E3G solution, 2ml of ovalbumin (10mg/ml) was added and this was incubated for 2.5hrs at room temperature with constant mixing. The conjugate was then dialysed for 16hrs against 1L of phosphate buffered saline containing 0.1% sodium azide.

Preparation of BSA- mimotope conjugate

Bovine serum albumin (BSA, 10mg, Sigma) was dissolved in 3 ml of conjugation buffer (sodium hydrogen carbonate buffer, 0.1M, pH 8.4) in a clean glass vial, mixing by suction and expulsion from a pipette tip. The mixture was left on a roller for one hour. The peptide mimotope as represented by SEQ ID NO:36 was dissolved in conjugation buffer. Peptide solution (1.0 ml @ 10mg/ml) and 10 μ l glutaraldehyde (high commercial grade, Sigma) were added to the BSA solution. The sealed vial was then agitated on a roller for

four hours at room temperature. The conjugate solution was then dialysed against sodium chloride (0.9%w/v) for 48h at 4°C.

Assays:

Peptide-mimotope-BSA conjugate (10 μ g/ml) and E3G-ovalbumin conjugate (3 μ g/ml) were dried separately into the wells of a microtitre plate (Becton Dickinson, CA, USA) from 50 μ l of solution in Phosphate Buffered Saline (PBS) overnight at room temperature. Wells were washed (4x PBS+0.01 %Tween20™), blocked for 1 hour with 0.1% casein in 100 μ l of PBS and washed 4x before use. 25 μ l aliquots of E3G in PBS (0-3 μ M) were added and incubated for 15mins before adding in 25 μ l of the MAb4155 anti-E3G antibody at 0.6 μ g/ml in PBS. The wells were incubated, with agitation, for 1 hour at room temperature. After washing 4x, 50 μ l of rabbit anti-mouse IgG-HRP conjugate (Dako, High Wycombe, UK) at 1:1000 dilution in PBS was added to each well and incubated for 1 hour at room temperature. After washing 4x, the wells were incubated in ABTS [2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid)] working substrate for 15 min (150 μ l/well). ABTS was prepared as a 0.033% (w/v) solution in 0.1M citrate phosphate buffer (pH 4.5) with 33% hydrogen peroxide (1 μ l/ml). Colour development was terminated by addition of 0.5M sulphuric acid (10 μ l/well) and was measured spectrophotometrically at 405 nm using a Milenia Kinetic Analyser™ (DPC, Llanberis, Wales).

Figure 1 shows the resulting assay curves. Both are typical for competitive immunoassays having midpoints in the micromolar to nanomolar range. Furthermore, the assay sensitivity using the mimotope-containing BSA-SEQ ID NO:37 is significantly greater than that obtained when using the epitope-containing E3G-ovalbumin.

The invention has been described in detail with particular reference to preferred embodiments thereof, but it will be understood that variations and modifications can be effected with the spirit and scope of the invention.